

confiderm® LDC

Textured Fingertips

Provides excellent tactile sensitivity and dexterity

Improved Conformability

Supports a comfortable fit for extended wear

Chemo Protection

Approved for use with chemotherapy drugs

Wrist Coverage

Coverage past the wrist helps protect against fluid run-off onto skin

Low Dermatitis Potential

Formulated to help reduce the risk of Type IV Dermatitis skin irritation for users with sensitive skin

- Tested for use with chemotherapy drugs per ASTM D6978
(See next page for specific drugs)

PRODUCT SPECIFICATIONS

Color	Blue
Cuff Style	Beaded Cuff
Fingertip Thickness Range	2 mil to 4 mil Range
Fingertip Thickness: Typ. Values	0.09 mm (3.6 mil)
Glove Exterior	Textured Fingertips
Glove Interior	Without Coating
Glove Length	Standard Cuff Length
Hand Compatibility	Ambidextrous
Length in Inches	9.6 Inch
Material	Nitrile
Protection Level	Chemo Tested
Sterility	Non-sterile
Not Made with Natural Rubber Latex	

**CHEMO
TESTED**

ORDERING INFORMATION

E1 #	MFR #	DESCRIPTION	SIZE	QTY
1281474	14-6N421C	Nitrile Exam Gloves, Chemo Tested, Non-sterile, Textured, Powder-Free	Small (S)	250/bx, 10bx/cs
1281475	14-6N441C	Nitrile Exam Gloves, Chemo Tested, Non-sterile, Textured, Powder-Free	Medium (M)	250/bx, 10bx/cs
1281476	14-6N461C	Nitrile Exam Gloves, Chemo Tested, Non-sterile, Textured, Powder-Free	Large (L)	250/bx, 10bx/cs
1281477	14-6N481C	Nitrile Exam Gloves, Chemo Tested, Non-Sterile, Textured, Powder-Free	X-Large (XL)	230/bx, 10 bx/cs

CHEMOTHERAPY DRUG PERMEATION RESISTANCE TESTED PER ASTM D6978

Chemotherapy Drug and Concentration Tested	Minimum Breakthrough Detection Time in Minutes
Carboplatin , 10 mg/mL (10,000 ppm)	>240
Carmustine (BCNU) (3,300 ppm or 3.3 mg/mL)	Do Not Use
Cisplatin (1,000 ppm or 1.0 mg/mL)	>240
Cyclophosphamide (Cytoxan) (20,000 ppm or 20.0 mg/mL)	>240
Dacarbazine (DTIC) (10,000 ppm or 10.0 mg/mL)	>240
Doxorubicin Hydrochloride (2,000 ppm or 2.0 mg/mL)	>240
Etoposide (Toposar) (20,000 ppm or 20.0 mg/mL)	>240
Fluorouracil (50,000 ppm or 50.0 mg/mL)	>240
Ifosfamide (50,000 ppm or 50.0 mg/mL)	>240
Methotrexate (25,000 ppm or 25.0 mg/mL)	>240
Mitomycin C (500 ppm or 0.5 mg/mL)	>240
Mitoxantrone (2,000 ppm or 2.0 mg/mL)	>240
Paclitaxel (Taxol) (6,000 ppm or 6.0 mg/mL)	>240
Thiotepa (10,000 ppm or 10.0 mg/mL)	Do Not Use
Vincristine Sulfate (1,000 ppm or 1.0 mg/mL)	>240

Warning: Do not use with Carmustine or Thiotepa.

Satisfaction Guaranteed

For complete details, please visit
mms.mckesson.com/mckesson-brands.

Never Compromise on Care™

With McKesson Brands, you'll never have to compromise. We deliver quality products at competitive prices, backed by our 100% satisfaction guarantee. We have decades of experience in the market, and we offer you support and work to help ensure you have the products you need, when you need them—making it even easier for you to provide the best care for your patients.

The product information contained in this document, including the product images and additional product materials, was collected from various supplier sources. All product claims and specifications are those of the product suppliers and have not been independently verified by McKesson Medical-Surgical or its affiliates ("McKesson"). McKesson is not responsible for errors or omissions in the product information.

The properties of a product may change or be inaccurate following the posting or printing of the product information in the document, either in the print or online version. Caution should be exercised when using or purchasing any products from McKesson's online or print documents by closely examining the product packaging and the labeling prior to use.

Due to product changes, information listed in this document is subject to change without notice. This information is placed solely for your convenience in ordering and McKesson disclaims all responsibility for its completeness and accuracy, whether or not the inaccuracy or incompleteness is due to fault or error by McKesson.